


Why use the SPICCT™?

SPICCT™ helps clinicians identify people with one or more general indicators of poor or deteriorating health and clinical signs of life-limiting conditions for assessment and care planning. SPICCT™ looks for changes in health status, burden of illness and increasing care needs. Integrate a holistic palliative care approach with best available treatment of underlying illnesses. Timely identification avoids harm and improves treatment and care of patients and families.

Using SPICCT™ to assess people’s needs and plan care.

- Unplanned **hospital admission**, more clinic **visits** or a **decline in health status**: review current care, treatment and medication; discuss future options; plan for managing further deterioration.
- **Poorly controlled symptoms**: review and optimise available treatment of underlying conditions; stop medicines/interventions/tests not of benefit; use effective palliative symptom control measures.
- People who are **increasingly dependent on others** due to deteriorating functional ability, physical frailty and/or mental health problems often need additional care and support.
- **Complex symptoms** or other patient/family **needs**; consider specialist palliative care review or involve another appropriate specialist or service.
- Assess **decision-making capacity**. Plan ahead if this will deteriorate. Record details of close family/friends, Power of Attorney or legal proxies. Involve in decision-making if capacity is impaired.
- Identify people who need proactive, **coordinated care in the community** from the primary care team and/or other community staff and services. Involve the local community. Support carers.
- Agree, record, share, and plan to review advance/anticipatory care plans (**ACP**); include plans for urgent/ emergency care and treatment if the person’s health deteriorates or care at home changes.

Talking about future care planning



- Talk about:
 - Benefits, harms and costs of hospital admission, outpatient visits, tests and treatments (e.g. IV antibiotics/fluids; surgery; cancer treatments, interventions for heart or kidney disease; tube feeding; oxygen/ventilation.
 - Treatments that will not work or have a poor outcome for this person. (eg. cardiopulmonary resuscitation)
 - Choosing legal proxy decision-makers in case the person’s decision-making capacity is lost in the future.
 - What a person would like; anything they do not want.
 - Help and support for family/ informal carers.

Tips on starting conversations about deteriorating health

- *I wish we had a treatment for... Could we talk about what **we can do** if that’s not possible?*
- *I am glad you feel better and **I hope** you will stay well, but **I am worried** that you could get ill again...*
- ***Can we talk** about how we manage **not knowing exactly** what will happen and when?*
- *If you got less well in the future, what would **be important** for you? What **would she say** about this?*
- ***Some people** want to talk about whether to go to hospital or be at home if they are seriously ill....*